

# Odległe wyniki leczenia drenażem wentylacyjnym przewlekłego wysiękowego zapalenia ucha środkowego u dzieci

Results of treatment with tympanostomy tubes in children with otitis  
media with effusion

*Beata Zielnik-Jurkiewicz, Olga Olszewska-Sosińska, Magdalena Rakowska*

## Summary

**Introduction:** Otitis media with effusion is a common problem in childhood. The tympanostomy tubes is a standard treatment of persistent otitis media with effusion in children which improves hearing level. However some pathological changes of the tympanic membrane and middle ear few years after treatment are observed. The aim of the study was prevalence of middle ear sequelae in children with persistent otitis media with effusion treated by tympanostomy tubes 4—10 years after treatment. **Material and methods:** The group of 61 patients aged between 1—13 with otitis media with effusion treated by tympanostomy tubes in Department of Otolaryngology Children's Hospital in Warsaw between 1994—2001 were controlled. 113 ears treated with tympanostomy tubes were examined 4—10 years after treatment. At the examination otomicroscope, pure-tone audiometry and tympanometry were used. **Results:** Out of the 113 ears after ventilation tubes insertion as a method of otitis media with effusion treatment in 83 (73%) developed one or more middle ear sequelae. Myringosclerosis in 47 ears (41,59%), segmental atrophy in 19 ears (16,81%), atrophy in 14 ears (12,38%), disfunction of Eustachian tube in 8 ears (7,07%), otitis media with effusion in 7 ears (6,19%), perforation of tympanic membrane in 6 ears (5,3%), chronic otitis media in 2 ears (1,76%), retraction pocket in 2 ears (1,76%) and tympanostomy tube in the middle ear cavity in 1 ear (0,88%) were found. No cases of cholesteatoma were found. In 22 ears (19,46%) the conductive hearing loss was greater than 20 dB. In 4 ears (3,5%) mixed hearing loss and in 3 ears (2,6%) receptive hearing loss was diagnosed. Tympanogram type A was in 86 ears (76,1%), tympanogram type B was in 11 ears (9,7%) and tympanogram type C was in 12 ears (12,6%). **Conclusions:** In children with otitis media with effusion treated with tympanostomy tubes the amount of sequelae increased with time. In children with otitis media with effusion who started treatment with tympanostomy tubes in age to 4 years old the amount of sequelae is less. The follow-up is recommended up to 8 years after the treatment.